

On the consistency strength of the proper forcing axiom

Matteo Viale, Christoph Weiß*

Abstract

In [21] the second author introduced combinatorial principles that characterize supercompactness for inaccessible cardinals. We prove that the proper forcing axiom PFA implies these principles hold for ω_2 . Using this, we argue to show that any of the known methods for forcing models of PFA from a large cardinal assumption requires a strongly compact cardinal. If one forces PFA using a proper forcing, then we get the optimal result that a supercompact cardinal is necessary.

1 Introduction

Since their introduction in the seventies supercompact cardinals played a central role in set theory. They have been a fundamental assumption to obtain many of the most interesting breakthroughs: Solovay's original proof that the singular cardinal hypothesis SCH holds eventually above a large cardinal, Silver's first proof of $\text{Con}(\neg\text{SCH})$, Woodin, Martin and Steel's first proofs of projective determinacy from large cardinals, Baumgartner's proof of the consistency of the proper forcing axiom PFA [2] and Foreman, Magidor, and Shelah's proof of the consistency of Martin's maximum MM [5] all relied on the assumption of the existence of a supercompact cardinal.

While some of these result have been shown to have considerably weaker consistency strength, the exact large cardinal strength of the forcing axioms PFA and MM is one of the major open problems in set theory. It is what we want to address in this paper.

*Parts of the results of this paper are from the second author's doctoral dissertation [22] written under the supervision of Dieter Donder, to whom the second author wishes to express his gratitude.

Forcing axioms play an important role in contemporary set theory. Historically they evolved from Martin's axiom, which was commonly used as the axiomatic counterpart to " $V = L$." The most prominent forcing axioms today are PFA as well as the stronger MM. Not only do they serve as a natural extension of ZFC, they also answer a plethora of questions undecidable in ZFC alone, from elementary questions like the size of the continuum to combinatorially complicated ones like the basis problem for uncountable linear orders [15]. Even problems originating from other fields of mathematics and apparently unrelated to set theory have been settled appealing to PFA. For example, Farah [3] recently proved the nonexistence of outer automorphisms of the Calkin algebra assuming PFA.

The consistency proofs of PFA and MM both start in a set theoretic universe in which there is a supercompact cardinal κ . They then collapse κ to ω_2 in such a way that in the resulting model PFA or MM holds, thus showing the consistency strength of these axioms is at most that of the existence of a supercompact cardinal.

An early result on PFA by Baumgartner [1] was that PFA implies the tree property on ω_2 , that is, PFA implies there are no ω_2 -Aronszajn trees. As a cardinal κ is weakly compact if and only if it is inaccessible and the tree property holds on κ , this can be seen as PFA showing the "weak compactness" of ω_2 , apart from its missing inaccessibility. This is an affirmation of the idea that collapsing a large cardinal to ω_2 is necessary to produce a model of PFA, and it actually implies the consistency strength of PFA is at least the existence of a weakly compact cardinal, for if the tree property holds on ω_2 , then ω_2 is weakly compact in L by [13].

This was the first insight that showed PFA possesses large cardinal strength, and many heuristic results indicate that supercompactness actually is the correct consistency strength of PFA and thus in particular also of MM. Still giving lower bounds for the consistency strength of PFA or MM is one major open problem today. While inner model theoretic methods were refined and enhanced tremendously over the last three decades, the best lower bounds they can establish today are still far below supercompactness [8].

In [21] the second author introduced combinatorial principles which do for strong compactness and supercompactness what the tree property does for weak compactness: A cardinal κ is strongly compact (supercompact) if and only if κ is inaccessible and $\text{TP}(\kappa)$ or, equivalently, $\text{SP}(\kappa)$ ($\text{ITP}(\kappa)$ or, equivalently, $\text{ISP}(\kappa)$) holds. We will show PFA implies $\text{ISP}(\omega_2)$, the strongest of the four principles. This, in the line of thought from above, says PFA shows ω_2 is, modulo inaccessibility, "supercompact."

Apart from the strong heuristic evidence this gives, by using arguments for pulling back these principles from generic extensions these characterizations actu-

ally allow us to show the following new kinds of theorems: If one forces a model of PFA using a reasonable forcing iteration that collapses a large cardinal to ω_2 , then this cardinal has to be strongly compact. If the iteration also is to be proper, then one needs a supercompact cardinal. Right now, such forcings are the only method both available and conceivable for producing models of PFA from large cardinal assumptions.

These results greatly improve on recent work by Neeman [16], who showed that if one starts with a ground model that satisfies certain fine structural properties and forces PFA by means of a proper forcing, then the ω_2 of the generic extension has to be a cardinal κ which is close to being κ^+ -supercompact in the ground model. (More precisely, in the ground model there is a Σ_1^2 -indescribable $[\kappa, \kappa^+]$ -gap.)

Notation

The notation used is mostly standard. For a regular cardinal δ , $\text{cof } \delta$ denotes the class of all ordinals of cofinality δ .

The phrases *for large enough* θ and *for sufficiently large* θ will be used for saying that there exists a θ' such that the sentence's proposition holds for all $\theta \geq \theta'$.

For an ordinal κ and a set X we let $P_\kappa X := \{x \subset X \mid |x| < \kappa\}$ and, if $\kappa \subset X$,

$$P'_\kappa X := \{x \in P_\kappa X \mid \kappa \cap x \in \text{Ord}, \langle x, \in \rangle < \langle X, \in \rangle\}.$$

For $x \in P_\kappa X$ we set $\kappa_x := \kappa \cap x$. For $f : P_\omega X \rightarrow P_\kappa X$ let $\text{Cl}_f := \{x \in P_\kappa X \mid \forall z \in P_\omega x \ f(z) \subset x\}$. Cl_f is club, and it is well known that for any club $C \subset P_\kappa X$ there is an $f : P_\omega X \rightarrow P_\kappa X$ such that $\text{Cl}_f \subset C$.

For sections 2 and 3, κ and λ are assumed to be cardinals, $\kappa \leq \lambda$, and κ is regular and uncountable.

Acknowledgments

The authors wish to express their gratitude to David Asperó, Sean Cox, Dieter Donder, Hiroshi Sakai, Ralf Schindler, and Boban Veličković for valuable comments and feedback on this research. They are indebted to Menachem Magidor for supplying them with the idea of the proof of Theorem 6.9, that is, Claim 6.9.1. They furthermore want to thank Mauro Di Nasso for an invitation to discuss this material at a one week workshop in Pisa.

2 The principles TP, SP, ITP, and ISP

We recall the necessary definitions from [21]. Let us call a sequence $\langle d_a \mid a \in P_\kappa\lambda \rangle$ a $P_\kappa\lambda$ -list if $d_a \subset a$ for all $a \in P_\kappa\lambda$.

Definition 2.1. Let $D = \langle d_a \mid a \in P_\kappa\lambda \rangle$ be a $P_\kappa\lambda$ -list.

- D is called *thin* if there is a club $C \subset P_\kappa\lambda$ such that $|\{d_a \cap c \mid c \subset a \in P_\kappa\lambda\}| < \kappa$ for every $c \in C$.
- D is called *slender* if for every sufficiently large θ there is a club $C \subset P_\kappa H_\theta$ such that $d_{M \cap \lambda} \cap b \in M$ for all $M \in C$ and all $b \in M \cap P_{\omega_1}\lambda$.

Note that if D is a thin list, then D is slender.

Definition 2.2. Let $D = \langle d_a \mid a \in P_\kappa\lambda \rangle$ be a $P_\kappa\lambda$ -list and $d \subset \lambda$.

- d is called a *cofinal branch of D* if for all $a \in P_\kappa\lambda$ there is $z_a \in P_\kappa\lambda$ such that $a \subset z_a$ and $d \cap a = d_{z_a} \cap a$.
- d is called an *ineffable branch of D* if there is a stationary set $S \subset P_\kappa\lambda$ such that $d \cap a = d_a$ for all $a \in S$.

Definition 2.3. • $\text{TP}(\kappa, \lambda)$ holds if every thin $P_\kappa\lambda$ -list has a cofinal branch.

- $\text{SP}(\kappa, \lambda)$ holds if every slender $P_\kappa\lambda$ -list has a cofinal branch.
- $\text{ITP}(\kappa, \lambda)$ holds if every thin $P_\kappa\lambda$ -list has an ineffable branch.
- $\text{ISP}(\kappa, \lambda)$ holds if every slender $P_\kappa\lambda$ -list has an ineffable branch.

We let $\text{TP}(\kappa)$ abbreviate the statement that $\text{TP}(\kappa, \lambda)$ holds for all $\lambda \geq \kappa$, and similarly for the other principles.

These definitions admit different ways of defining strong compactness and supercompactness.

Theorem 2.4. *Suppose κ is inaccessible. Then κ is strongly compact if and only if $\text{TP}(\kappa)$ holds.*

Theorem 2.5. *Suppose κ is inaccessible. Then κ is supercompact if and only if $\text{ITP}(\kappa)$ holds.*

Unlike other characterizations however, by [21] the principles ITP and ISP also make sense for small cardinals.

There exist ideals and filters naturally associated to the principles ITP and ISP.

Definition 2.6. Let $A \subset P_\kappa \lambda$ and let $D = \langle d_a \mid a \in P_\kappa \lambda \rangle$ be a $P_\kappa \lambda$ -list. D is called *A-effable* if for every $S \subset A$ that is stationary in $P_\kappa \lambda$ there are $a, b \in S$ such that $a \subset b$ and $d_a \neq d_b \cap a$. D is called *effable* if it is $P_\kappa \lambda$ -effable.

Definition 2.7. We let

$$I_{\text{IT}}[\kappa, \lambda] := \{A \subset P_\kappa \lambda \mid \text{there exists a thin } A\text{-effable } P_\kappa \lambda\text{-list}\},$$

$$I_{\text{IS}}[\kappa, \lambda] := \{A \subset P_\kappa \lambda \mid \text{there exists a slender } A\text{-effable } P_\kappa \lambda\text{-list}\}.$$

By $F_{\text{IT}}[\kappa, \lambda]$ and $F_{\text{IS}}[\kappa, \lambda]$ we denote the filters associated to $I_{\text{IT}}[\kappa, \lambda]$ and $I_{\text{IS}}[\kappa, \lambda]$ respectively.

The ideals $I_{\text{IT}}[\kappa, \lambda]$ and $I_{\text{IS}}[\kappa, \lambda]$ are normal ideals on $P_\kappa \lambda$ by [21].

3 Guessing models

We now introduce the concept of a *guessing model* which gives an alternative presentation of the principle ISP.

Definition 3.1. Let $M < H_\theta$ for some large enough θ .

- A set d is called *M-approximated* if $d \cap b \in M$ for all $b \in M \cap P_{\omega_1} M$.
- A set d is called *M-guessed* if there is an $e \in M$ such that $d \cap M = e \cap M$.

M is called *z-guessing* if every M -approximated $d \subset z$ is M -guessed. M is called *guessing* if for all $z \in M$, M is z -guessing.

Note that since for every $z \in M$ there is a bijection $f : z \rightarrow \rho$ in M for some ordinal ρ , it holds that M is guessing if and only if M is ρ -guessing for all $\rho \in M$. Also note that since M cannot be $\sup(M \cap \text{Ord})$ -guessing, any ordinal ρ such that M is ρ -guessing has to be bounded by $\sup(M \cap \text{Ord})$.

Define

$$\mathcal{G}_\kappa^z X := \{M \in P'_\kappa X \mid M \text{ is } z\text{-guessing}\},$$

$$\mathcal{G}_\kappa X := \{M \in P'_\kappa X \mid M \text{ is guessing}\}.$$

Proposition 3.2. *If $\text{ISP}(\kappa, |H_\theta|)$ holds, then $\mathcal{G}_\kappa H_\theta$ is stationary.*

Proof. By working with a bijection $f : |H_\theta| \rightarrow H_\theta$, it is obvious that we can apply $\text{ISP}(\kappa, |H_\theta|)$ to the set $P_\kappa H_\theta$ directly.

Suppose to the contrary that there is a club $C \subset P'_\kappa H_\theta$ such that every $M \in C$ is not guessing, that is, there is $z_M \in M$ and $d_M \subset z_M$ that is M -approximated but not M -guessed. Then also $d_M \cap M$ is M -approximated but not M -guessed, so we may assume $d_M \subset M$. Consider the list $D := \langle d_M \mid M \in C \rangle$.

Then D is slender, for let θ' be large enough and let $C' := \{M' \in P_\kappa H_{\theta'} \mid M' \cap H_\theta \in C\}$. C' is club in $P_\kappa H_{\theta'}$, and if $M' \in C'$ and $b \in P_{\omega_1} H_{\theta'} \cap M'$, then $b \in M' \cap H_\theta$, so $d_{M' \cap H_\theta} \cap b \in M' \cap H_\theta \subset M'$.

By $\text{ISP}(\kappa, |H_\theta|)$, there is an ineffable branch d for the list D . Let $S := \{M \in C \mid d_M = d \cap M\}$. S is stationary, and we may assume $z_M = z$ for some fixed z and all $M \in S$. This means $d \subset z$. As $Pz \subset H_\theta$, there is an $M \in S$ such that $d \in M$. But then d_M is M -guessed, a contradiction. \square

Proposition 3.3. *Let θ be sufficiently large and $M \in P'_\kappa H_\theta$ be a λ -guessing model such that $\lambda^+ \in M$. Then $\text{ISP}(\kappa, \lambda)$ holds.*

Proof. Since $M < H_\theta$ it is enough to show that $M \models \text{ISP}(\kappa, \lambda)$. So pick a slender list $D = \langle d_a \mid a \in P_\kappa \lambda \rangle \in M$. Notice that the slenderness of D is witnessed by a club $C' \subset P_\kappa H_{\lambda^+}$ which is in M . Then $M \cap H_{\lambda^+} \in C'$, so $d_{M \cap \lambda} \cap b \in M$ for all $b \in M \cap P_{\omega_1} \lambda$. This means $d_{M \cap \lambda}$ is an M -approximated subset of M . So since M is a λ -guessing model, there is an $e \in M$ such that $e \cap M = d_{M \cap \lambda}$.

Let $S := \{a \in P_\kappa \lambda \mid d_a = e \cap a\}$. Then $S \in M$. To see S is stationary, let $C \in M$ be a club in $P_\kappa \lambda$. Then $M \cap \lambda \in C \cap S$, so $H_\theta \models C \cap S \neq \emptyset$, so it also holds in M . \square

Notice that we cannot literally say that $F_{\text{IS}}[\kappa, H_\theta]$ is the club filter restricted to $\mathcal{G}_\kappa H_\theta$: There might be a slender list $\langle d_M \mid M \in S \rangle$ indexed by some stationary set $S \subset \mathcal{G}_\kappa H_\theta$ that does not have an ineffable branch. For such a list we necessarily have that $d_M \not\subset z$ for all $z \in M$ and all $M \in S$. Still the following holds.

Proposition 3.4. *$I_{\text{IS}}[\kappa, X]$ is contained in the projection of the nonstationary ideal restricted to $\mathcal{G}_\kappa^X H_\theta$ onto X for any regular θ such that $X \in H_\theta$.*

Proof. Assume to the contrary that there is an $S \in I_{\text{IS}}[\kappa, X]$ such that $S^* := \{M \in \mathcal{G}_\kappa^X H_\theta \mid M \cap X \in S\}$ is stationary. Pick a slender list $D = \langle d_a \mid a \in S \rangle$ witnessing that $S \in I_{\text{IS}}[\kappa, X]$. Let C be a club subset of $P_\kappa H_\theta$ witnessing that D is slender. Pick $M \in S^* \cap C$ such that $D \in M$. Then $d_{M \cap X}$ is an M -approximated subset of X as

$M \in \mathcal{C}$. Thus $d_{M \cap X} = e \cap M$ for some $e \in M$ since M is X -guessing. As in the proof of Proposition 3.3 it follows that e is an ineffable branch for D , contradicting the fact that D witnesses $S \in I_{\text{IS}}[\kappa, X]$. \square

4 Implications under PFA

In this section, we are going to show PFA implies $\text{ISP}(\omega_2)$.

The following lemma is due to Woodin [23, Proof of Theorem 2.53]. Recall that $G \subset \mathbb{P}$ is said to be M -generic if G is a filter on \mathbb{P} and $G \cap D \cap M \neq \emptyset$ for all $D \in M$ that are dense in \mathbb{P} .

Lemma 4.1. *Let \mathbb{P} be a proper forcing, and let θ be sufficiently large. Then PFA implies*

$$\{M \in P_{\omega_2}H_\theta \mid \exists G \subset \mathbb{P} \text{ } G \text{ is } M\text{-generic}\}$$

is stationary in $P_{\omega_2}H_\theta$.

Definition 4.2. Let T be a tree and B be a set of cofinal branches of T . A function $g : B \rightarrow T$ is called *Baumgartner function* if g is injective and for all $b, b' \in B$ it holds that

1. $g(b) \in b$,
2. $g(b) < g(b') \rightarrow g(b') \notin b$.

The following lemma is due to Baumgartner, see [1].

Lemma 4.3. *Let T be a tree and B be a set cofinal branches of T . Suppose $\kappa := \text{ht}(T)$ is regular and $|B| \leq \kappa$. Then there is a Baumgartner function $g : B \rightarrow T$.*

Proof. Let $\langle b_\alpha : \alpha < \mu \rangle$ enumerate B , with $\mu \leq \kappa$. Recursively define g by $g(b_\alpha) := \min(b_\alpha - \bigcup \{b_\beta : \beta < \alpha\})$. This can be done since κ is regular. Suppose $g(b_\alpha) < g(b_{\alpha'})$ for some $\alpha, \alpha' < \mu$. Then $g(b_{\alpha'}) \in b_{\alpha'}$, so $g(b_\alpha) \in b_{\alpha'}$, so $\alpha < \alpha'$ and thus $g(b_{\alpha'}) \notin b_\alpha$. \square

Recall that a tree T is said to *not split at limit levels* if for all $t, t' \in T$ such that $\text{ht } t = \text{ht } t'$ is a limit ordinal and $\{s \in T : s < t\} = \{s \in T : s < t'\}$ it follows that $t = t'$.

Lemma 4.4. *Let T be a tree that does not split at limit levels and suppose B is a set of cofinal branches of T . Suppose $g : B \rightarrow T$ is a Baumgartner function. Suppose $\langle \alpha_\nu : \nu < \omega_1 \rangle$ is continuous and increasing. Let $\alpha := \sup_{\nu < \omega_1} \alpha_\nu$ and $t \in T_\alpha$. Suppose that for all $\nu < \omega_1$ there is $b_\nu \in B$ such that $g(b_\nu) < t \upharpoonright \alpha_\nu \in b_\nu$. Then there is a stationary $S \subset \omega_1$ such that $b_\nu = b_{\nu'}$ for all $\nu, \nu' \in S$. In particular there is an $s < t$ such that $t \in g^{-1}(s)$.*

Proof. For $\nu < \omega_1$ let $r(\nu) := \min\{\rho < \nu \mid \text{ht } g(b_\nu) < \alpha_\rho\}$. Then r is regressive and thus constant on a stationary set $S \subset \omega_1$. As g is a Baumgartner function, this implies g is constant on the set $\{b_\nu \mid \nu \in S\}$. But g is injective, so $b_\nu = b_{\nu'}$ for $\nu, \nu' \in S$. \square

Definition 4.5. Let $V \subset W$ be a pair of transitive models of ZFC.

- (V, W) satisfies the μ -covering property if the class $P_\mu^V V$ is cofinal in $P_\mu^W V$, that is, for every $x \in W$ with $x \subset V$ and $|x| < \mu$ there is $z \in P_\mu^V V$ such that $x \subset z$.
- (V, W) satisfies the μ -approximation property if for all $x \in W$, $x \subset V$, it holds that if $x \cap z \in V$ for all $z \in P_\mu^V V$, then $x \in V$.

A forcing \mathbb{P} is said to satisfy the μ -covering property or the μ -approximation property if for every V -generic $G \subset \mathbb{P}$ the pair $(V, V[G])$ satisfies the μ -covering property or the μ -approximation property respectively.

These properties have been introduced and extensively studied by Hamkins, see for example [7].

The following lemma is the essential argument in the proof of Theorem 4.8. Extracting it has the advantage that it can be applied to a wider class of different forcings, so that it can yield more information about the nature of the guessing models and $I_{\mathbb{S}}[\omega_2, \lambda]$.

Lemma 4.6. *Let θ be sufficiently large. Assume \mathbb{P} satisfies the ω_1 -covering and the ω_1 -approximation properties and collapses 2^λ to ω_1 . Then in $V^{\mathbb{P}}$ there is a ccc forcing $\dot{\mathbb{Q}}$ and some $w \in H_\theta$ such that*

$$\{M \in P'_{\omega_2} H_\theta \mid w \in M, \exists G \subset \mathbb{P} * \dot{\mathbb{Q}} \text{ } G \text{ is } M\text{-generic}\} \subset \mathcal{G}_\kappa^\lambda H_\theta,$$

and every such M is internally unbounded, that is, $M \cap P_{\omega_1} M$ is cofinal in $P_{\omega_1} M$.

Proof. Let $B := {}^{\omega_1}2$.

Work in $V^{\mathbb{P}}$. Let $\dot{c} : \omega_1 \rightarrow P_{\omega_1}\lambda$ be continuous and cofinal. As \mathbb{P} satisfies the ω_1 -covering property, we may assume that $\dot{c}(\alpha + 1) \in V$ for all $\alpha < \omega_1$. Define

$$\dot{T} := \{h \upharpoonright \dot{c}(\alpha) \mid h \in B, \alpha < \omega_1\}$$

As \mathbb{P} satisfies the ω_1 -approximation property, we have that B is the set of cofinal branches through \dot{T} .

Since $|B| = \omega_1$, we can apply Lemma 4.3 and get a Baumgartner function $\dot{g} : B \rightarrow \dot{T}$. Let $\dot{l} : \omega_1 \rightarrow B$ be a bijection. Let

$$\begin{aligned} \dot{T}^0 &:= \{t \in \dot{T} : \exists b \in B \dot{g}(b) < t \in b\}, \\ \dot{T}^1 &:= \dot{T} - \dot{T}^0. \end{aligned}$$

Note that \dot{T}^1 does not have cofinal branches. Thus there is a ccc forcing $\dot{\mathbb{Q}}$ that specializes \dot{T}^1 with a specialization map \dot{f} .

Now work in V . Let $w \in H_\theta$ contain all the relevant information, and let $M \in P'_{\omega_2}H_\theta$ be such that $w \in M$ and there is an M -generic $G_0 * G_1 \subset \mathbb{P} * \dot{\mathbb{Q}}$.

By the usual density arguments, $c := \dot{c}^{G_0} : \omega_1 \rightarrow P_{\omega_1}(M \cap \lambda)$ is continuous and cofinal and $c(\alpha + 1) \in M$ for all $\alpha < \omega_1$. Therefore M is internally unbounded. We let $g := \dot{g}^{G_0}$, $T := \dot{T}^{G_0}$, $T^0 := (\dot{T}^0)^{G_0}$, $T^1 := (\dot{T}^1)^{G_0}$, $l := \dot{l}^G$, and $f := \dot{f}^{G_0 * G_1}$. Define $B \upharpoonright M := \{h \upharpoonright M \mid h \in B \cap M\}$. Then we can use the facts that $G_0 * G_1$ is an M -generic filter and that $V^{\mathbb{P}} \models \text{rng } \dot{l} = B$ to argue that

- $l : \omega_1 \rightarrow B \cap M$ is bijective,
- $T = \bigcup \{h \upharpoonright c(\alpha) \mid h \in B \cap M, \alpha < \omega_1\}$,
- $g : B \upharpoonright M \rightarrow T$ is a Baumgartner function,¹
- $T = T^0 \cup T^1$,
- $f : T^1 \rightarrow \omega$ is a specialization map.

Claim 4.6.1. $B \upharpoonright M$ is the set of uncountable branches of T .

¹Here we naturally identify $\text{dom } g = B \cap M$ with $B \upharpoonright M$, which is a set of uncountable branches of T .

Proof. It is clear that $B \upharpoonright M$ is included in the set of uncountable branches of T . For the other inclusion, observe that if h is a branch through T , then h must be a branch through T_0 since the specialization map f witnesses that T_1 cannot have uncountable branches. This means that $h \upharpoonright c(\alpha) \in T_0$ for eventually all α . So for each such α there is a unique $b_\alpha \in B \upharpoonright M$ such that $g(b_\alpha) \subset h \upharpoonright c(\alpha) \subset b_\alpha$. Thus for eventually all $\alpha < \omega_1$ we have $\text{dom } g(b_\alpha) = c(\beta_\alpha)$ for some $\beta_\alpha < \alpha$, and we may assume that there is a $\beta < \omega_1$ such that $\beta_\alpha = \beta$ for stationarily many $\alpha < \omega_1$. Hence if α is such that $\beta_\alpha = \beta$, then $h = b_\alpha \in B \upharpoonright M$. \dashv

Claim 4.6.2. $t \in B \upharpoonright M$ if and only if t is the characteristic function of $d \cap M$ for some M -approximated $d \subset \lambda$.

Proof. If $t \in B \upharpoonright M$, then $t = h \upharpoonright M$ for some $h \in B \cap M$, and h is the characteristic function of some $d \in M \cap P\lambda$.

For the other direction pick an M -approximated $d \subset \lambda$, and let t be the characteristic function of $d \cap M$. We claim that t is a branch through T and thus in $B \upharpoonright M$ by Claim 4.6.1. To see this observe that $c(\alpha + 1) \in M$ for all $\alpha < \omega_1$, so that $t \upharpoonright c(\alpha + 1)$ is the characteristic function of $d \cap c(\alpha + 1)$, which is in M since d is M -approximated. Thus $t \upharpoonright c(\alpha + 1) \in T$. \dashv

To see M is λ -guessing, let $d \subset \lambda$ be M -approximated. Then by Claim 4.6.2 the characteristic function t of $d \cap M$ is in $B \upharpoonright M$. So there is $h \in B \cap M$ such that $t = h \upharpoonright M$. Let $e \in M$ be such that h is its characteristic function. Then $e \cap M = d \cap M$, and we are done. \square

To apply Lemma 4.6, we need an appropriate forcing. The simplest and earliest example comes from [13]. We let \mathbb{C} denote the forcing for adding a Cohen real. See [11] for a proof of the following theorem.

Theorem 4.7. *Let $\gamma \geq \omega_1$. Then the forcing $\mathbb{C} * \text{Coll}(\omega_1, \gamma)$ is proper and satisfies the ω_1 -approximation property.*

Theorem 4.8. *PFA implies $\text{ISP}(\omega_2)$ holds.*

Proof. Let θ be large enough, $\lambda \geq \omega_2$, and $\mathbb{P} := \mathbb{C} * \text{Coll}(\omega_1, 2^\lambda)$. Then \mathbb{P} is proper and satisfies the ω_1 -approximation property by Theorem 4.7. Thus by Lemmas 4.1 and 4.6 the set $\mathcal{G}_{\omega_2}^\lambda H_\theta$ is stationary in $P_{\omega_2} H_\theta$. Therefore by Proposition 3.3 we can conclude that $\text{ISP}(\omega_2, \lambda)$ holds. \square

Krueger [10; 12] has shown there is a great variety of forcings $\dot{\mathbb{P}}$ living in $V^{\mathbb{C}}$ such that $\mathbb{C} * \dot{\mathbb{P}}$ has the ω_1 -approximation and the ω_1 -covering properties. These forcings can be used to show that under PFA, there are stationarily many guessing models that are internally club. As guessing models are not internally approachable, this gives another separation of the properties internally club and internally approachable. Under MM, one can use these forcings to show there are stationarily many guessing models that are internally unbounded but not internally stationary and also stationarily many that are internally stationary but not internally club, see also [20].

Strullu [18] has shown the principle $\text{ITP}(\omega_2)$ follows from $\text{MRP} + \text{MA}$, where MRP is the mapping reflection principle introduced by Moore [14].

It is furthermore worth noting that unlike $\text{ISP}(\omega_2)$, the principle $\text{ITP}(\omega_2)$ can already be proved by applying PFA to a forcing of the form σ -closed $*$ ccc, see [22].

The next corollary is originally independently due to Foreman and Todorćević, see [9].

Corollary 4.9. *PFA implies the approachability property fails for ω_1 , that is, $\omega_2 \notin I[\omega_2]$, where $I[\omega_2]$ denotes the approachability ideal on ω_2 .*

Proof. It is not hard to see that $I[\omega_2] \subset I_{\text{IS}}[\omega_2, \omega_2]$. □

The failure of various square principles under PFA is originally due to Todorćević and Magidor, see [19] and [17, Theorem 6.3]. See [21] for the notation used in Corollary 4.10.

Corollary 4.10. *Suppose PFA holds and cf $\lambda \geq \omega_2$. Then $\neg \square_{\text{cof}(\omega_1)}(\omega_2, \lambda)$.*

Proof. This follow from Theorem 4.8 and [21, Theorem 4.2]. □

5 An interlude on forcing

Definition 5.1. Let \mathbb{P} be a forcing. We say \mathbb{P} is a *standard iteration of length κ* if

- (i) \mathbb{P} is the direct limit of an iteration $\langle \mathbb{P}_\alpha \mid \alpha < \kappa \rangle$ that takes direct limits stationarily often,
- (ii) \mathbb{P}_α has size less than κ for all $\alpha < \kappa$.

It is a classical result that the μ -cc is preserved by iterations of length μ of posets of size less than μ that take direct limits stationarily often. So the following lemma does not come as a surprise but nonetheless has not been observed so far.

Lemma 5.2. *Let \mathbb{P} be a standard iteration of length κ . Then \mathbb{P} is κ -cc and satisfies the κ -approximation property.*

Proof. Let \mathbb{P} be the direct limit of $\langle \mathbb{P}_\alpha \mid \alpha < \kappa \rangle$. It suffices to verify the κ -approximation property for subsets of ordinals. The proof is by induction on $\lambda \geq \kappa$.

We start with the proof of the base case $\lambda = \kappa$. We need to show that if $p \in \mathbb{P}$ and $\dot{h} \in V^{\mathbb{P}}$ are such that $p \Vdash_{\mathbb{P}} \dot{h} \in {}^\kappa 2$ and $p \Vdash_{\mathbb{P}} \forall \alpha < \kappa \dot{h} \upharpoonright \alpha \in V$, then $p \Vdash_{\mathbb{P}} \dot{h} \in V$. So assume to the contrary there is $\bar{p} \leq p$ such that $\bar{p} \Vdash_{\mathbb{P}} \dot{h} \notin V$.

Let $P = \{p_\xi \mid \xi < \kappa\}$ and let C_0 be the club of all $\alpha < \kappa$ such that $\bigcup \{\mathbb{P}_\xi \mid \xi < \alpha\} = \{p_\xi \mid \xi < \alpha\}$. Define $S := \{\alpha < \kappa \mid \mathbb{P}_\alpha \text{ is direct limit}\}$. S is stationary by assumption, and if $\alpha \in S \cap C_0$, then $\mathbb{P}_\alpha = \{p_\xi \mid \xi < \alpha\}$.

For $\xi < \kappa$ let $A_\xi \subset \mathbb{P}$ be a maximal antichain below \bar{p} that decides the value of $\dot{h}(\xi)$. Then $C := \{\alpha \in C_0 \mid \forall \xi < \alpha A_\xi \subset \mathbb{P}_\alpha\}$ is club. For $\alpha \in C$ let

$$\dot{h}_\alpha := \{\langle (\xi, i), p \rangle \mid \xi < \alpha, p \in \mathbb{P}_\alpha, p \Vdash_{\mathbb{P}} \dot{h}(\xi) = i\}.$$

Then $\dot{h}_\alpha \in V^{\mathbb{P}_\alpha}$ and $\bar{p} \Vdash_{\mathbb{P}} \dot{h}_\alpha \in {}^\alpha 2$.

Claim 5.2.1. $\bar{p} \Vdash_{\mathbb{P}} \dot{h} \upharpoonright \alpha = \dot{h}_\alpha$ for all $\alpha \in C$.

Proof. Suppose to the contrary that for some $\alpha \in C$ there are $q \leq \bar{p}$ and $\xi < \alpha$ such that $q \Vdash_{\mathbb{P}} \dot{h}(\xi) \neq \dot{h}_\alpha(\xi)$. Let $r \in A_\xi$ be compatible with q . Then $r \Vdash_{\mathbb{P}} \dot{h}(\xi) = i$ for some $i < 2$. But as $A_\xi \subset \mathbb{P}_\alpha$, this also means $r \Vdash_{\mathbb{P}} \dot{h}_\alpha(\xi) = i$, contradicting its compatibility with q . \dashv

Claim 5.2.2. $\bar{p} \Vdash_{\mathbb{P}_\alpha} \dot{h}_\alpha \in V$ for all $\alpha \in C$.

Proof. Assume towards a contradiction that some for some $q \leq \bar{p}$ and $\alpha \in C$ we have $q \Vdash_{\mathbb{P}_\alpha} \dot{h}_\alpha \notin V$. Then for each $g \in {}^\alpha 2$ there is a maximal antichain A_g among the conditions in \mathbb{P}_α below q such that for any element $r \in A_g$, there is $\xi_r < \alpha$ such that $r \Vdash_{\mathbb{P}_\alpha} \dot{h}_\alpha(\xi_r) \neq g(\xi_r)$. This means that any $\langle (\xi_r, i), p \rangle \in \dot{h}_\alpha$ such that p is compatible with r is such that $g(\xi_r) \neq i$. This in turn means that $r \Vdash_{\mathbb{P}} \dot{h}_\alpha(\xi_r) \neq g(\xi_r)$ for any $r \in A_g$ and for any $g \in {}^\alpha 2$.

Since a maximal antichain in \mathbb{P}_α is also a maximal antichain in \mathbb{P} , this implies that $q \Vdash_{\mathbb{P}} \dot{h}_\alpha \notin V$, which is impossible by Claim 5.2.1. \dashv

For $\alpha \in S \cap C_0$ by Claim 5.2.2 $\bar{p} \Vdash_{\mathbb{P}_\alpha} \dot{h}_\alpha \in V$, so there are $p_\xi \in \mathbb{P}_\alpha$, $p_\xi \leq \bar{p}$, and $g_\alpha \in {}^\alpha 2$ such that $p_\xi \Vdash_{\mathbb{P}_\alpha} \dot{h}_\alpha = g_\alpha$. Since $\alpha \in S \cap C_0$, we have $\xi < \alpha$, so for some

stationary $S_0 \subset S \cap C_0$ we may assume ξ is fixed. But then $p_\xi \Vdash_{\mathbb{P}_\alpha} \dot{h} \upharpoonright \alpha = \dot{h}_\alpha = g_\alpha$ for all $\alpha \in S_0$, so that $p_\xi \Vdash_{\mathbb{P}} \dot{h} = \bigcup_{\alpha \in S_0} \dot{h}_\alpha = \bigcup_{\alpha \in S_0} g_\alpha \in V$, contradicting $p_\xi \leq \bar{p}$.

Now we prove the lemma for $\lambda > \kappa$, assuming it has been shown for all $\gamma < \lambda$. Let $p \in \mathbb{P}$ and $\dot{h} \in V^{\mathbb{P}}$ be such that $p \Vdash_{\mathbb{P}} \dot{h} \in {}^\lambda 2$ and $p \Vdash_{\mathbb{P}} \forall z \in P_\kappa^V V \dot{h} \upharpoonright z \in V$.

First suppose cf $\lambda > \kappa$. By the induction hypothesis we know that $p \Vdash_{\mathbb{P}} \forall \gamma < \lambda \dot{h} \upharpoonright \gamma \in V$. For every $\gamma < \lambda$ there is $\alpha_\gamma < \kappa$ and $g_\gamma \in {}^\gamma 2$ such that $p_{\alpha_\gamma} < p$ and $p_{\alpha_\gamma} \Vdash \dot{h} \upharpoonright \gamma = g_\gamma$. Thus there is an unbounded $U \subset \lambda$ such that $\alpha_\gamma = \alpha_{\gamma'}$ for all $\gamma, \gamma' \in U$, so that for $\gamma \in U$ we have $p_{\alpha_\gamma} \Vdash \dot{h} = \bigcup_{\gamma \in U} g_\gamma \in V$.

If cf $\lambda \leq \kappa$, let $U \subset \lambda$ be cofinal of order type cf λ , and set

$$T := \{g \in {}^{<\lambda} 2 \mid \exists q \leq p \exists \gamma \in U q \Vdash_{\mathbb{P}} \dot{h} \upharpoonright \gamma = g\}.$$

Then T , ordered by end extension, is a tree of height cf λ . As \mathbb{P} is κ -cc, all levels of T have size less than κ . Let X be a set of size at most κ such that for every pair of incompatible elements $g, g' \in T$ there is $\alpha \in X$ such that $g(\alpha) \neq g'(\alpha)$. By the induction hypothesis we have $p \Vdash_{\mathbb{P}} \dot{h} \upharpoonright X \in V$. But $p \Vdash_{\mathbb{P}} \dot{h} = \bigcup \{g \in T \mid g \upharpoonright X = \dot{h} \upharpoonright X\}$, so that $p \Vdash_{\mathbb{P}} \dot{h} \in V$. \square

6 The principles TP and ITP in generic extensions

Lemma 6.1. *Let $V \subset W$ be a pair of models of ZFC that satisfies the κ -covering property, and suppose κ is inaccessible in V . Suppose $D = \langle d_a \mid a \in P_\kappa^W \lambda \rangle$ is a $P_\kappa^W \lambda$ -list such that for every $a \in P_\kappa^W \lambda$ there is $z_a \in V$ such that $d_a = z_a \cap a$. Then D is thin.*

Proof. Work in W . Let $c \in P_\kappa \lambda$. By the κ -covering property there is $\bar{c} \in P_\kappa^V \lambda$ such that $c \subset \bar{c}$. Also we have $\{d_a \cap c \mid c \subset a \in P_\kappa^W \lambda\} = \{z_a \cap \bar{c} \cap c \mid c \subset a \in P_\kappa^W \lambda\} \subset \{z \cap c \mid z \in P^V \bar{c}\}$. But the latter set has cardinality less than κ since κ is inaccessible in V . \square

Proposition 6.2. *Let $V \subset W$ be a pair of models of ZFC that satisfies the κ -covering and the κ -approximation properties, and suppose κ is inaccessible in V . Then*

$$I_{\text{IT}}^V[\kappa, \lambda] \subset I_{\text{IT}}^W[\kappa, \lambda].$$

Proof. Work in W . For $A \in I_{\text{IT}}^V[\kappa, \lambda]$ let $\langle d_a \mid a \in P_\kappa^V \lambda \rangle \in V$ be A -effable in V .

Then by Lemma 6.1 $\langle d_a \mid a \in P_\kappa \lambda \rangle$ is thin, where $d_a := \emptyset$ for $a \notin V$.

Suppose $\langle d_a \mid a \in P_\kappa \lambda \rangle$ were not A -effable. Let $S \subset A$ be stationary and $d \subset \lambda$ such that $d_x = d \cap x$ for all $x \in S$. Suppose $d \notin V$. Then, by κ -approximation property, there is a $z \in P_\kappa^V \lambda$ such that $d \cap z \notin V$. But for $x \in S$ with $z \subset x$ we have $d \cap z = d \cap x \cap z = d_x \cap z \in V$, a contradiction. Therefore $d \in V$, and $S \subset \bar{S} := \{x \in P_\kappa^V \lambda \mid d_x = d \cap x\} \in V$. Since $\langle d_a \mid a \in P_\kappa^V \lambda \rangle \in V$ is A -effable in V , \bar{S} is not stationary in V . So there exists $C \in V$, $C \subset P_\kappa^V \lambda$ club in V such that $C \cap \bar{S} = \emptyset$. Let $f : P_\omega \lambda \rightarrow P_\kappa \lambda$ be in V such that $\text{Cl}_f^V \subset C$. But then, by the stationarity of S , there is an $x \in S$ such that $x \in \text{Cl}_f$, so that $x \in C \cap \bar{S}$, a contradiction. \square

Theorem 6.3. *Let $V \subset W$ be a pair of models of ZFC that satisfies the κ -covering property and the τ -approximation property for some $\tau < \kappa$, and suppose κ is inaccessible in V . Then*

$$P_\kappa^W \lambda - P_\kappa^V \lambda \in I_{\text{IT}}^W[\kappa, \lambda],$$

which furthermore implies

$$F_{\text{IT}}^V[\kappa, \lambda] \subset F_{\text{IT}}^W[\kappa, \lambda].$$

So in particular, if $W \models \text{ITP}(\kappa, \lambda)$, then $V \models \text{ITP}(\kappa, \lambda)$.

Proof. Work in W . Let $B := P_\kappa \lambda - P_\kappa^V \lambda$. For $x \in B$ let $a_x \in P_\tau^V \lambda$ be such that $x \cap a_x \notin V$, which exists by the τ -approximation property. Put $d_x := a_x \cap x$. For $x \in P_\kappa \lambda - B$, let $d_x := \emptyset$. Then $\langle d_x \mid x \in P_\kappa \lambda \rangle$ is thin by Lemma 6.1.

Suppose $\langle d_x \mid x \in P_\kappa \lambda \rangle$ were not B -effable. Then there are $d \subset \lambda$ and $U \subset B$ be such that U is cofinal and $d_x = d \cap x$ for all $x \in U$. Define a \subset -increasing sequence $\langle x_\alpha \mid \alpha < \tau^+ \rangle$ with $x_\alpha \in U$ for all $\alpha < \tau^+$ and a sequence $\langle e_\alpha \mid \alpha < \tau^+ \rangle$ such that $x_\alpha \subset e_\alpha$ and $e_\alpha \in P_\kappa^V \lambda$ for all $\alpha < \tau^+$ as follows. Let $\beta < \tau^+$ and suppose $\langle x_\alpha \mid \alpha < \beta \rangle$ and $\langle e_\alpha \mid \alpha < \beta \rangle$ have been defined. Let $x_\beta \in U$ be such that $\bigcup_{\alpha < \beta} (x_\alpha \cup a_\alpha \cup e_\alpha) \subset x_\beta$, and let $e_\beta \in P_\kappa^V \lambda$ be such that $x_\beta \subset e_\beta$, which exists by the κ -covering property.

Then $\langle d_{x_\alpha} \mid \alpha < \tau^+ \rangle$ is \subset -increasing as $d_{x_\alpha} = d \cap x_\alpha$ for all $\alpha < \tau^+$, and since $|d_{x_\alpha}| < \tau$ for all $\alpha < \tau^+$, there is $\gamma < \tau^+$ such that $d_{x_\alpha} = d_{x_{\alpha'}}$ for all $\alpha, \alpha' \in [\gamma, \tau^+)$. But then $a_{x_{\gamma+1}} \cap e_\gamma \subset a_{x_{\gamma+1}} \cap x_{\gamma+1} = d_{x_{\gamma+1}} = d_{x_\gamma} \subset e_\gamma$ and $d_{x_{\gamma+1}} \subset a_{x_{\gamma+1}}$, so that $d_{x_\gamma} = a_{x_{\gamma+1}} \cap e_\gamma \in V$, a contradiction.

To see $F_{\text{IT}}^V[\kappa, \lambda] \subset F_{\text{IT}}^W[\kappa, \lambda]$, let $A \in F_{\text{IT}}^V[\kappa, \lambda]$. Then $P_\kappa^V \lambda - A \in I_{\text{IT}}^V[\kappa, \lambda]$, so by Proposition 6.2 $P_\kappa^V \lambda - A \in I_{\text{IT}}^W[\kappa, \lambda]$. Thus $P_\kappa^W \lambda - A = (P_\kappa^W \lambda - P_\kappa^V \lambda) \cup (P_\kappa^V \lambda - A) \in I_{\text{IT}}^W[\kappa, \lambda]$, which means $A \in F_{\text{IT}}^W[\kappa, \lambda]$. \square

Note that by [6, Theorem 1.1] the set $P_\kappa^W \lambda - P_\kappa^V \lambda$ in Theorem 6.3 is stationary for $\lambda \geq \kappa^+$ if there is a real in $W - V$. We will now weaken the assumption that (V, W) satisfies the τ -approximation property for some $\tau < \kappa$ to the κ -approximation property, so that this kind of argument can be exploited for a wider range of forcing constructions.

Theorem 6.4. *Let $V \subset W$ be a pair of models of ZFC that satisfies the κ -covering and the κ -approximation properties, and suppose κ is inaccessible in V . If $W \models \text{TP}(\kappa, \lambda)$, then $V \models \text{TP}(\kappa, \lambda)$.*

Proof. In V , let $D = \langle d_a \mid a \in P_\kappa \lambda \rangle$ be a $P_\kappa \lambda$ -list.

Now work in W . For every $a \in P_\kappa \lambda$ let, by the κ -covering property, $z_a \in P_\kappa^V \lambda$ be such that $a \subset z_a$. Define a $P_\kappa \lambda$ -list $E = \langle e_a \mid a \in P_\kappa \lambda \rangle$ by $e_a := d_{z_a} \cap a$. Then E is thin by Lemma 6.1.

Thus by $\text{TP}(\kappa, \lambda)$ there is a cofinal branch d for E . So for all $y \in P_\kappa \lambda$ there is $a \in P_\kappa \lambda$, $y \subset a$, such that $e_a \cap y = d \cap y$. In particular

$$d \cap y = e_a \cap y = d_{z_a} \cap a \cap y = d_{z_a} \cap y.$$

Thus if $y \in P_\kappa^V \lambda$, then $d \cap y \in V$, so that $d \in V$ by the κ -approximation property. This means $d \in V$. But d is also a cofinal branch for D in V . \square

Corollary 6.5. *Let \mathbb{P} be a standard iteration of length κ and suppose κ is inaccessible. If \mathbb{P} forces $\text{TP}(\kappa)$, then κ is strongly compact.*

Proof. This follows directly from Lemma 5.2 and Theorem 6.4. \square

Notice that, together with Theorem 4.8, Corollary 6.5 implies the following remarkable corollary.

Corollary 6.6. *Suppose κ is inaccessible and PFA is forced by a standard iteration of length κ that collapses κ to ω_2 . Then κ is strongly compact.*

Corollary 6.6 says that any of the known methods for producing a model of PFA from a large cardinal assumption requires at least a strongly compact cardinal. This can be improved to the optimal result if we require the iteration for forcing PFA to be proper. For this purpose we introduce an ad-hoc definition.

Definition 6.7. Let $V \subset W$ be a pair of models of ZFC that satisfies the κ -covering and the κ -approximation properties, and suppose κ is inaccessible in V . We say $M \in (P_\kappa^V H_\theta^V)^W$ is V -guessing if for all $z \in M$ and all $d \in P^V z$ there is an $e \in M$ such that $d \cap M = e \cap M$.

The following two propositions should be seen as analogs of Propositions 3.2 and 3.3.

Proposition 6.8. *Let $V \subset W$ be a pair of models of ZFC that satisfies the κ -covering and the κ -approximation properties, and suppose κ is inaccessible in V . Assume $W \models \text{ITP}(\kappa, |H_\theta^V|)$ for some large enough θ . Then in W the set*

$$\{M \in P'_\kappa H_\theta^V \mid M \text{ is } V\text{-guessing and closed under countable suprema}\}$$

*is stationary.*²

Proof. Work in W . By [21, Theorem 3.5], we have that the set of all $M \in P'_\kappa H_\theta^V$ that are closed under countable suprema belongs to $F_{\text{IT}}[\kappa, H_\theta^V]$. Assume that there were a set $A \notin I_{\text{IT}}[\kappa, H_\theta^V]$ such that for all $M \in A$ there is $z_M \in M$ and $d_M \in P^V z_M$ such that $d_M \cap M \neq e \cap M$ for all $e \in M$. Then $D := \langle d_M \cap M \mid M \in A \rangle$ is thin by Lemma 6.1. Thus by $\text{ITP}(\kappa, |H_\theta^V|)$ there is an ineffable branch d for D , and by the κ -approximation property we have $d \in V$. Let $S := \{M \in A \mid d_M \cap M = d \cap M\}$. Then $S \in V$ is stationary, and we may assume $z_M = z$ for some $z \in H_\theta^V$ and all $M \in S$. As $P^V z \subset H_\theta^V$ and $d \subset z$, there is an $M \in S$ such that $d \in M$, a contradiction. \square

Theorem 6.9. *Let $V \subset W$ be a pair of models of ZFC that satisfies the κ -covering and the κ -approximation properties. Let κ be inaccessible in V and λ be regular in W . Suppose that for all $\gamma < \kappa$ and every $S \subset \text{cof}(\omega) \cap \gamma$ in V it holds that $V \models$ “ S is stationary in γ ” if and only if $W \models$ “ S is stationary in γ .” Let θ be large enough. Suppose $M \in (P'_\kappa H_\theta^V)^W$ is a V -guessing model closed under countable suprema such that $\lambda \in M$. Then $M \cap \lambda \in V$ and $V \models \text{ITP}(\kappa, \lambda)$.*

Proof. Let $\langle S_\alpha \mid \alpha < \lambda \rangle \in M$ be a partition of $\text{cof}(\omega) \cap \lambda$ into sets stationary in V . Let $\lambda_M := \sup(M \cap \lambda)$.

Claim 6.9.1. *It holds that*

$$M \cap \lambda = \{\delta < \lambda \mid V \models S_\delta \text{ is stationary in } \lambda_M\} \in V.$$

Proof. For one direction, let δ be such that $V \models$ “ S_δ is stationary in λ_M .” Notice that $\text{cf}^V \lambda_M < \kappa$, so $W \models$ “ S_δ is stationary in λ_M .” As M is closed under countable suprema, we get that $S_\delta \cap M \neq \emptyset$. Thus if $\beta \in S_\delta \cap M$, then δ is definable in M as the α for which $\beta \in S_\alpha$, so that $\delta \in M$.

²However, it need not be a subset of V .

For the other direction, let $\delta \in M \cap \lambda$ and let $C \in V$ be club in λ_M . As $C \cap \lambda \in M$ and M is V -guessing, $C \cap M = e \cap M$ for some $e \in M$. Since $C \cap M$ is closed under countable suprema, $M \models$ “ e is closed under countable suprema.” Thus $M \models e \cap S_\delta \neq \emptyset$, which proves $C \cap S_\delta \neq \emptyset$ as $e \cap S_\delta \cap M \subset C \cap S_\delta$. \dashv

Now to argue that $V \models \text{ITP}(\kappa, \lambda)$, it is enough to check that $H_\theta^V \models \text{ITP}(\kappa, \lambda)$. Since $M < H_\theta^V$, it in turn suffices to verify $M \models \text{ITP}(\kappa, \lambda)$. So let $D \in M$ be a $P_\kappa^V \lambda$ -list. Since M is V -guessing, $d_{M \cap \lambda} \in V$, and $d_{M \cap \lambda} \subset \lambda \in M$, we get that $d_{M \cap \lambda} = e \cap M$ for some $e \in M$. Then $M \models$ “ e is an ineffable branch for D .” \square

Corollary 6.10. *Let \mathbb{P} be a proper standard iteration of length κ and suppose κ is inaccessible. If \mathbb{P} forces $\text{ITP}(\kappa)$, then κ is supercompact.*

Proof. This follows from Lemma 5.2, Proposition 6.8, and Theorem 6.9. \square

Under the additional premise of properness, Corollary 6.10 implies the following strongest possible version of Corollary 6.6.

Corollary 6.11. *Suppose κ is inaccessible and PFA is forced by a proper standard iteration of length κ that collapses κ to ω_2 . Then κ is supercompact.*

It should be noted that Sakai has pointed out a serious obstruction in removing the assumption of \mathbb{P} being proper in Corollary 6.11.

Theorem 6.12 (Sakai, 2010). *Let κ be a supercompact cardinal, $\theta > \kappa$ be sufficiently large, and suppose there is a Woodin cardinal $\mu > \theta$. Suppose W is the standard semiproper forcing extension such that $W \models \text{MM} + \kappa = \omega_2$. Then in W it holds that for every stationary preserving forcing \mathbb{P} the set*

$$\{M \in P_{\omega_2} H_\theta \mid \exists G \subset \mathbb{P} \text{ } G \text{ is } M\text{-generic, } M \cap \omega_3 \notin V\}$$

is stationary in $P_{\omega_2} H_\theta$.

In the setting of Theorem 6.12, if one carries out the proof of Theorem 4.8 in W , one gets that $P_\kappa^W \lambda - P_\kappa^V \lambda \notin I_{\text{IT}}^W[\kappa, \lambda]$ for λ such that $\kappa < \lambda$ and $2^\lambda < \theta$. This should be contrasted with Theorem 6.3.

7 Conclusion

There are several open problems which the results presented suggest. The most appealing deals with the construction of an inner model in which ω_2 has an arbitrary degree of supercompactness starting from a universe of sets in which MM holds. It seems plausible to conjecture that if $\text{ISP}(\kappa)$ holds, then for each λ there is a simply definable transitive class in which κ is λ -supercompact. Such a line of thought has already been pursued by Foreman [4], where he proved that a certain strong form of Chang's conjecture for a small cardinal κ implies that there is an X such that κ is huge in $L[X]$. It has yet to be understood to what extent Foreman's ideas can be applied to the results of this paper; a key issue in this context appears to be a thorough study of the properties of guessing models and of the ideals $I_{\text{IS}}[\omega_2, \lambda]$ in models of MM.

We also expect that many of the known consequences of PFA and supercompactness might be obtained directly from the principle ISP. Examples are given in [21], where it is shown that $\text{ITP}(\omega_2)$ implies the failure of some of the weakest forms of square incompatible with PFA, and in [20], where, using properties of guessing models, a new proof that PFA implies SCH is provided. On the other hand we conjecture that $\text{ISP}(\omega_2)$ does not decide the size of the continuum.

References

- [1] J. E. Baumgartner, *Applications of the proper forcing axiom*, Handbook of set-theoretic topology, North-Holland, Amsterdam, 1984, pp. 913–959. MR 776640
- [2] K. J. Devlin, *The Yorkshireman's guide to proper forcing*, Surveys in set theory, London Math. Soc. Lecture Note Ser., vol. 87, Cambridge Univ. Press, Cambridge, 1983, pp. 60–115. MR 823776
- [3] I. Farah, *All automorphisms of the Calkin algebra are inner*, Ann. of Math. (2), forthcoming.
- [4] M. Foreman, *Smoke and mirrors: combinatorial properties of small cardinals equiconsistent with huge cardinals*, Adv. Math. **222** (2009), no. 2, 565–595. MR 2538021
- [5] M. Foreman, M. Magidor, and S. Shelah, *Martin's maximum, saturated ideals*,

- and nonregular ultrafilters. I*, Ann. of Math. (2) **127** (1988), no. 1, 1–47. MR 924672
- [6] M. Gitik, *Nonsplitting subset of $\mathcal{P}_\kappa(\kappa^+)$* , J. Symbolic Logic **50** (1985), no. 4, 881–894 (1986). MR 820120
- [7] J. D. Hamkins, *Gap forcing*, Israel J. Math. **125** (2001), 237–252. MR 1853813
- [8] R. Jensen, E. Schimmerling, R. Schindler, and J. Steel, *Stacking mice*, J. Symbolic Logic **74** (2009), no. 1, 315–335. MR 2499432
- [9] B. König and Y. Yoshinobu, *Fragments of Martin’s maximum in generic extensions*, MLQ Math. Log. Q. **50** (2004), no. 3, 297–302. MR 2050172
- [10] J. Krueger, *Internally club and approachable*, Adv. Math. **213** (2007), no. 2, 734–740. MR 2332607
- [11] ———, *A general Mitchell style iteration*, MLQ Math. Log. Q. **54** (2008), no. 6, 641–651. MR 2472470
- [12] ———, *Internal approachability and reflection*, J. Math. Log. **8** (2008), no. 1, 23–39. MR 2674000
- [13] W. J. Mitchell, *Aronszajn trees and the independence of the transfer property*, Ann. Math. Logic **5** (1972/73), 21–46. MR 313057
- [14] J. T. Moore, *Set mapping reflection*, J. Math. Log. **5** (2005), no. 1, 87–97. MR 2151584
- [15] ———, *A five element basis for the uncountable linear orders*, Ann. of Math. (2) **163** (2006), no. 2, 669–688. MR 2199228
- [16] I. Neeman, *Hierarchies of forcing axioms. II*, J. Symbolic Logic **73** (2008), no. 2, 522–542. MR 2414463
- [17] E. Schimmerling, *Combinatorial principles in the core model for one Woodin cardinal*, Ann. Pure Appl. Logic **74** (1995), no. 2, 153–201. MR 1342358
- [18] R. Strullu, *MRP, tree properties and square principles*, forthcoming.

- [19] S. Todorčević, *A note on the proper forcing axiom*, *Axiomatic set theory* (Boulder, Colo., 1983), *Contemp. Math.*, vol. 31, Amer. Math. Soc., Providence, RI, 1984, pp. 209–218. MR 763902
- [20] M. Viale, *On the notion of guessing model*, forthcoming.
- [21] C. Weiß, *The combinatorial essence of supercompactness*, forthcoming.
- [22] ———, *Subtle and ineffable tree properties*, Ph.D. thesis, Ludwig Maximilians Universität München, 2010.
- [23] W. H. Woodin, *The axiom of determinacy, forcing axioms, and the nonstationary ideal*, *de Gruyter Series in Logic and its Applications*, vol. 1, Walter de Gruyter & Co., Berlin, 1999. MR 1713438